



MALATHION	GROUP	1B	INSECTICIDE
-----------	-------	----	-------------

FIFRA SECTION 24(c) SPECIAL LOCAL NEED (SLN) LABEL FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF GEORGIA

MALATHION 57 EC

EPA Reg. No. 34704-108

SLN No. GA-250002

FOR INDOOR USE ONLY FOR CONTROL OF BED BUGS IN COMMERCIAL POULTRY BREEDER HOUSES

This label is valid until December 31, 2030, or until otherwise amended,
withdrawn, canceled, or suspended.

Active Ingredient:	% BY WT.
Malathion*	57.00%
Other Ingredients	43.00%
Total	100.0%

*O, O-Dimethyl phosphorodithioate of diethyl mercaptosuccinate

Keep Out of Reach of Children CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalles. (If you do not understand this label, find someone to explain it to you in detail).

ENVIRONMENTAL HAZARDS

Follow all applicable directions, restrictions, and precautions in the Environmental Hazards section of the label affixed to the main container of Malathion 57 EC (EPA Reg. No. 34704-108).

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This SLN label and the label affixed to the main container of Malathion 57 EC must be in the possession of the user at the time of application.

Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the label affixed to the main container of Malathion 57 EC.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

POLLINATOR PROTECTION: Protect pollinating insects by following label directions intended to minimize drift and reduce risk to these organisms.

Target Area	Target Pest	Use Rate	Application Instructions
Commercial Poultry Breeder Houses	Bed Bugs	1.0 gallon of Malathion 57 EC per 25.0 gallons of diluent Apply: 3 gal/1000 sq feet (0.6 lbs. ai/1000 sq ft.)	<p>Prior to application, all breeder houses are required to be free of birds and treated after clean-out. Do not allow residues to directly contact litter, sweepings, or debris during application.</p> <p>Use a hand-held sprayer or power sprayer from a non-ULV application system.</p> <p>Make a thorough application when applying the finished application.</p>

SPECIFIC USE INSTRUCTIONS:

Only for indoor use in poultry breeder houses with birds removed during the clean-out process. Do not allow residues to directly contact poultry litter, sweepings, or debris during application.

This product may be applied by broadcast and/or crack and crevice treatment for use as a residual spray on walls, floors, ceilings, and other items in the breeder house that will not be affected by the spray.

The maximum application rate is 0.6 lbs. ai/1000 sq. ft. (1.0 pint of Malathion 57 EC Insecticide / 1000 sq. ft.). When mixed at a rate of 1 gal of this product to 25 gallons of water, apply mixture at 3 gals of finished solution / 1000 sq ft.

Restricted Entry Interval – 12 hours

BUFFER ZONE FOR APPLICATION NEAR WATER

Do not apply within 25 feet of aquatic habitats (such as, but not limited to, lakes, reservoirs, rivers, permanent streams, wetlands, or natural ponds, estuaries, and commercial fishponds).

ADDITIONAL WATER PROTECTIONS STATEMENTS for non-agricultural uses

- Do not apply directly to or allow the product to enter sewers or storm drains, or to any area like a drain or gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur.
- Do not apply directly to poultry litter, sweepings, or debris.
- Do not apply directly to impervious horizontal surfaces such as sidewalks, driveways, and patios except as a spot or crack-and-crevice treatment.
- Do not apply to vertical surfaces directly above pervious or impervious surfaces that drain into ditches, storm drains, gutters, or surface waters.
- Do not apply to the point of runoff.